NASA Satellite Trace Gas Products for Air Quality Applications

ARSET Webinar: April, 2014

ARSET

Applied Remote Sensing Education and Training



Focus: Three Remote Sensing Trace Gas Products

1. Tropospheric Column NO₂

 From OMI (Ozone Monitoring Instrument) on-board Aura satellite (http://aura.gsfc.nasa.gov)

2. Boundary Layer SO2


From Aura OMI

3. Total Column and profile measurements of CO

 From MOPITT (Measurements of Pollution in the Troposphere) on-board the Terra satellite (http://www2.acd.ucar.edu/mopitt)



Aura/OMI NO₂



OMI-derived tropospheric NO₂ columns track NO₂ pollution near the surface, which is an important precursor of urban smog.



Level 1 – Raw

Highest Resolution, Difficult

Level 2 - Orbit

Level 3 - Gridded



Lower Resolution, Easy



OMI Tropospheric Column NO₂ for AQ appl.

- Two Separate Algorithms
 - NASA/GSFC "Standard Product"
 - Level 3 data 0.25x0.25 degree gridded
 - Data Short name = OMNO2d
 - 2. KNMI (Royal Netherlands Meteorological Institute) DOMINO (Near-Real Time) Product
 - Level 2 data and images available from KNMI and TEMIS.
 - Data Short Name = OMDOMINO
 - http://www.temis.nl/airpollution/no2.html
 - Includes data/images from European Satellite instruments: GOME, GOME-2, SCIAMACHY



GSFC Standard Product for AQ Applications

Data Set Short Name = OMNO2d

Product Level = 3

Begin Date = October 1, 2004

Resolution = 0.25°lon x 0.25°lat

Version = 003


Cloud-screened best observation

Production Frequency: Daily

Granule (File) Coverage: 15 orbits

File Size (Approx): 2.5 MB

Data Format = HDF (hierarchical data format)



Data are publicly available here:

http://disc.sci.gsfc.nasa.gov/Aura/data-holdings/OMI/omno2d v003.shtml




Ozone Monitoring Instrument




Important information regarding OMI


Almost 50% data loss since 2008 (row anomaly effect)


Affects O₃, SO₂, and to some extent NO₂ OMI Products





Examples of OMI NO₂ Applications









OMI DOMINO Product: Level 2 only

Direct observations of megacity emissions and lifetimes of nitrogen oxides






Mean NO₂ tropospheric columns in the Middle East from OMI measurements during 2005-2009 for calm (w<2 m/s) conditions.

Mean NO₂ column densities around Riyadh for different wind conditions. Arrows indicate the mean of the respective ECMWF winds.

By analyzing the downwind patterns of NO₂ for different wind conditions, megacity NOx emissions and lifetimes from OMI can be determined.

OMNO₂ Standard Product shows declining trends over the Eastern US ~10%/year




David Streets and Zifeng Lu

Decision and Information Sciences Division

Argonne National Laboratory

Uses OMI NO₂ tropospheric columns to constrain models to identify NOx emission sources.



OMI NO₂ Data Access

GES-DISC (Goddard Earth Science Data and Information Services Center)

- http://disc.sci.gsfc.nasa.gov/Aura/data-holdings/OMI/omno2d_v003.shtml
- Level 2, L2G, Level 3 (OMNO2d)
- HDF and NetCDF
- Documentation

AVDC (Aura Validation Data Center)


- http://avdc.gsfc.nasa.gov
- Level 3 (OMNO2d)
- HDF, ASCII, KML formats
- Daily and monthly data and Images

GIOVANNI – an interactive visualization tool

- http://disc.sci.gsfc.nasa.gov/giovanni
- L2G, Level 3 (OMNO2d)
- HDF, ASCII, KML for Google Earth
- Subsetting available




http://disc.sci.gsfc.nasa.gov/Aura/data-holdings/OMI/omno2d_v003.shtml




-Aura Validation Data Center (AVDC)

NASA GES-DISC - OMNO2d, Level 3, 0.25 Degree Product


http://disc.sci.gsfc.nasa.gov/Aura/data-holdings/OMI/omno2d_v003.shtml







File Name Convention





OMI NO₂ Data Access

GES-DISC (Goddard Earth Science Data and Information Services Center)

- http://disc.sci.gsfc.nasa.gov/Aura/data-holdings/OMI/omno2d_v003.shtml
- Level 2, L2G, Level 3 (OMNO2d)
- HDF and NetCDF
- Documentation


AVDC (Aura Validation Data Center)


- http://avdc.gsfc.nasa.gov
- Level 3 (OMNO2d)
- HDF, ASCII, KML formats
- Daily and monthly data and Images

GIOVANNI – an interactive visualization tool

- http://disc.sci.gsfc.nasa.gov/giovanni
- L2G, Level 3 (OMNO2d)
- HDF, ASCII, KML for Google Earth
- Subsetting available











AVDC - Aura Validation Data Center http://avdc.gsfc.nasa.gov/index.php?site=705441739








KML – filename "EOS-Aura-OMI-OMNO2-L3_2011_01_NO2TropCS30.kml"





Tropospheric NO2 CS30 vertical column densities (VCD)







Google Earth image of OMI Tropospheric NO2 for











OMI NO₂ Data Access

GES-DISC (Goddard Earth Science Data and Information Services Center)

- http://disc.sci.gsfc.nasa.gov/Aura/data-holdings/OMI/omno2d_v003.shtml
- Level 2, L2G, Level 3 (OMNO2d)
- HDF and NetCDF
- Documentation

AVDC (Aura Validation Data Center)

- http://avdc.gsfc.nasa.gov
- Level 3 (OMNO2d)
- HDF, ASCII, KML formats
- Daily and monthly data and Images

GIOVANNI – an interactive visualization tool

- http://disc.sci.gsfc.nasa.gov/giovanni
- L2G, Level 3 (OMNO2d)
- HDF, ASCII, KML for Google Earth
- Subsetting available












http://gdata1.sci.gsfc.nasa.gov/daac-bin/G3/gui.cgi?instance_id=omi








NASA

Aura/OMI SO₂



A case study of the 2009 eruption of Mt. Redoubt, Alaska, US The dispersion of volcanic ash was monitored using OMI column SO_2 and CALIPSO backscatter data.

OMI SO2 Gridded Product Summary



- The OMI data product file contains 4 estimates of the total SO₂ column in Dobson Units.
- These correspond to 4 a-priori (climatological) vertical profiles for the SO₂ used in the retrieval algorithm.
- The 4 vertical profiles represent typical SO₂ vertical distributions for 4 SO₂ source regimes:

SO ₂ Product	Level	Data Short Name	Sensitivity	Use
PBL SO ₂	L3, 0.25°x0.25°	OMSO2e	< 1 km	Fossil fuel, industry
TRL SO ₂ (Lower Troposphere)	L2G, 0.25°x0.25°	OMSO2G	~ 3 km	Industry outflow
TRM SO ₂ (Mid-Tropopshere)	L2G, 0.25°x0.25°	OMSO2G	5 - 10 km	optimized for volcanic degassing with vents at ~5km altitude or above and emissions from effusive eruptions.
STL SO ₂ (Lower Stratosphere)	L2G, 0.25°x0.25°	OMSO2G	15 – 20 km	intended for use with explosive volcanic eruptions

Note: OMSO2e 'best' product: Screens anomalous values due to cloudiness, high terrain and instrumental effects (row anomaly)

. L2G data are NOT screened.

OMI AQ SO₂ product in the boundary Layer

Data Set Short Name = OMSO2e

Product Level = 3

Begin Date = October 1, 2004

Resolution = 0.25° lon x 0.25° lat

Version = 003


Cloud-screened best measurement

Production Frequency: Daily

Granule (File) Coverage: 15 orbits

File Size(Approx): 5 MB

Contains **best** pixel data, screened for OMI row anomaly, clouds, and other data quality flags.



Aqua MODIS visible image of the Nabro (Eritrea) eruption on June 13, 2011 and the SO2 plume overlaid.



Examples of OMI SO₂ Applications



Aura/OMI measurements show distinct enhancements in NO₂ and SO₂ amounts over the Canadian oil sands













■ Surface Mining Operations



OMI SO₂ Fort McMurray 0.1 0.2 0.3 0.4






Data from the Ozone Monitoring Instrument (OMI) reveal distinct enhancements in nitrogen dioxide (NO₂) and sulfur dioxide (SO₂) over a region of surface mining in the Canadian oil sands. The oil sands are located in the north-east corner of the province of Alberta, Canada and contain the second largest reserve (after Saudi Arabia) of oil globally. Shown along the top are maps over a portion of the oil sands where extensive surface mining occurs. These maps cover the same area: a NASA LandSat image from 2009 indicating vegetation removal in region of surface mining (left), OMI mean NO₂ (middle), and OMI mean SO₂ (right). OMI data were averaged over the period 2005-2010. White values indicate background levels, increasing through yelloworange, with brown-black the largest. Maximum values clearly correspond to the location of the industrial activity. The time graph shows that NO₂ has been increasing at a rate of 10%/year over this period. Maximum values are comparable to those seen over medium-to-large North American power plants.

These results make use of improved OMI mapping techniques to resolve detail down to 5 miles. These are the first satellite-based results of pollution over the oil sands.

Aura/OMI SO₂




http://earthobservatory.nasa.gov/IOTD/view.php?id=82626



Mean SO₂ values for 2005-2007

Mean SO₂ values for 2008-2010



The Ozone Monitoring Instrument (OMI) data confirm a substantial reduction in sulfur dioxide (SO_2) values around the largest US coal power plants as a result of the implementation of SO_2 pollution control measures. The figure shows average SO_2 values measured by OMI on the NASA Aura spacecraft for the periods 2005-2007 and 2008-2010 over the Eastern US where the majority of large SO_2 sources are located. Scientists use this information to identify anthropogenic sources of SO_2 and to estimate their emission rates. The greatest values are in violet; the lowest in green. Yellow to violet colors correspond to statistically significant enhancements in SO_2 pollution in the vicinity of largest SO_2 emitting coal-burning power plants indicated by the black dots.

power plants

Previous use of space-based SO₂ retrievals has been limited to monitoring plumes from volcanic eruptions and detecting anthropogenic emissions from large source regions as in China. A new spatial filtration technique allows detection of individual pollution sources in Canada and US.

Fioletov, V., et al., (2011), Geophysical Research Letters

Data/Images Access

Global Sulfur Dioxide Monitoring Homepage

- http://so2.gsfc.nasa.gov/
- Images, Documentation, Publications and Links

GES-DISC (Goddard Earth Science Data and Information Services Center)

- Level 2, L2G, Level 3 (OMSO2e)
- HDF and NetCDF
- Documentation

GIOVANNI – An interactive visualization tool

- L2G, Level 3 (OMSO2e)
- HDF, ASCII, KML for Google Earth
- Subsetting available





National Aeronautics and Space Administration Goddard Space Flight Center http://


Search SED Site

GO

Flight Projects | Sciences and Exploration

Atmospheric Chemistry and Dynamics Laboratory (Code 614)

Global Sulfur Dioxide Monitoring Home Page





National Aeronautics and Space Administration Goddard Space Flight Center

http://so2.gsfc.nasa.gov

Search SED Site Flight Projects | Sciences and Exploration

Near-real time SO₂ maps over Europe only

ric Chemistry and Dynamics Laboratory (Code 614)

Sulfur Dioxide Monitoring Home Page

Past SO₂ images Documentation Publications Personnel Links

Volcanic Hazards Project

test SO₂ eruption alerts NOAA-NESDIS. SACS_BIRA. IASI-ULB.

SO₂ Near Real-Time Images: Real-Time (15 mins). NOAA Near Real-Time (3 hours). AIRS NRT. NASA NRT. SACS-BIRA


NOAA Level 2 volcanic SO₂ images and archived data (various formats, specific sites)

MEaSUREs project:

ges (1979-2005) | AID images (2003-2004) | OMI images (2004-present) | OMPS images (May 2012-present)

aily (OM/OMPS) Images of SO₂ (click on a highlighted rectangle)

canic regions, orange = daily pollution regions, yellow = long-term pollution images



Data/Images Access

Global Sulfur Dioxide Monitoring Homepage

- http://so2.gsfc.nasa.gov/
- Images, Documentation, Publications and Links

GES-DISC (Goddard Earth Science Data and Information Services Center)

- http://disc.sci.gsfc.nasa.gov/Aura/data-holdings/OMI/omso2e_v003.shtml
- Level 2, L2G, <u>Level 3 (OMSO2e)</u>
- HDF and NetCDF
- Documentation

GIOVANNI – An interactive visualization tool


- L2G, Level 3 (OMSO2e)
- HDF, ASCII, KML for Google Earth
- Subsetting available

Worldview - Images






http://disc.sci.gsfc.nasa.gov/Aura/data-holdings/OMI/omso2e_v003.shtml






http://disc.sci.gsfc.nasa.gov/Aura/data-holdings/OMI/omso2e_v003.shtml







- + Feedback
- + FAQ

NASA Search Results

(Number of files found may not be entirely accurate)


Page: 1




NASA Privacy Policy and Important Notices

⁺ Contact Us

File Name Convention







Data/Images Access

Global Sulfur Dioxide Monitoring Homepage

- http://so2.gsfc.nasa.gov/
- Images, Documentation, Publications and Links

GES-DISC (Goddard Earth Science Data and Information Services Center)

- http://disc.sci.gsfc.nasa.gov/Aura/data-holdings/OMI/omso2e_v003.shtml
- Level 2, L2G, <u>Level 3 (OMSO2e)</u>
- HDF and NetCDF
- Documentation

GIOVANNI – An interactive visualization tool

- http://disc.sci.gsfc.nasa.gov/giovanni
- L2G, <u>Level 3 (OMSO2e)</u>
- HDF, ASCII, KML for Google Earth
- Subsetting available



Worldview - Images

GES DISC Home

Data Services

Science Portals

Mission Portals

Analyze Data with Giovanni


Search for Data with Mirador

Simple Subset Wizard

Data Cookbook

More...

OVANNI - The Bridge Between Data and Science





» OVERVIEW

- + What is Giovanni?
- + Who Uses Giovanni?
- + Giovanni Parameters
- + Giovanni Plot Types
- + How to Use Giovanni
- + How to Acknowledge Giovanni
- + Acknowledgements

Giovanni - Interactive Visualization and Analysis

You are here: GES DISC Home » Giovanni - Interactive Visualization and Analysis

Contributors: tonyr, rchowdhury

Giovanni - Interactive Visualization an<mark>d</mark> Analysis - GES <mark>D</mark>ISC: Go<mark>dd</mark>ard Earth <mark>Sci</mark>ences, <mark>D</mark>ata an<mark>d</mark> Information Services Center

Giovanni-4 Now Available

New! Please try out Giovanni-4, the next generation of Giovanni, with dramatically improved performance and interactive plotting and mapping. (Currently, only select Aerosols, Hydrology and Turbulent Flux data are available in Giovanni-4, with more on the way.)

Additional Features

- + News
- + Users Manual
- + Publications
- + Newsletters
- + Feedback
- + FAQ

Giovanni Portals

Giovanni Parameter List

- Atmospheric Portals (Scroll cown to view complete list)
 - Agua/AIRS Global: Monthly
 - Terra and Aqua MODIS: Daily
 - Terra and Aqua MODIS: Monthly
 - Aura OMI Level 3
 - Aura OMI Level 2G
 - Aura Microwave Limb Sounder (MLS)
 - Aura High Resolution Dynamics Limb Sounder (HIRDLS)
 - Aura Tropospheric Emission Spectrometer (TES)
 - Earth Probe and Nimbus-7 TOMS
 - Upper Atmosphere Research Satellite (UARS) Halogen Occultation Experiment (HALOE)

Application and Education Bortal (Sarall Hours to view complete list

GIOVANNI NEWS

Giovanni Image Hall of Fame issue of The Giovanni News is online

Jan 10, 2014

Several members of the GES DISC attend ESIP Federation Winter Meeting 2014

Jan 07, 2014

December 2013 AGU special issue of The Giovanni News is online

Dec 19, 2013

GES DISC participates in AGU Fall Meeting 2013

Dec 06, 2013

October-November 2013 issue of The Giovanni News is online

Nov 22, 2013

MODIS observes progressive development of air pollution crisis in China


Oct 25, 2013

Staff from the GES DISC participate in NSF EarthCube Workshop


Oct 21, 2013

Newest additions to Giovanni publications list

Sep 30, 2013




http://disc.sci.gsfc.nasa.gov/giovanni




http://disc.sci.gsfc.nasa.gov/giovanni O' S' S' O' O A Search DISC 1/4/2013 8:00:00 pm + G0 Ø 3 € 6 1/4/2013 8:00:00 pm Chandigarh Uttarakhand Himalaya Mountair Haryana/ Delhi/ दिल्लो Kathmandu 分面図 Uttar Pradesh Rajasthan OMSO2e,003 Vertical Column Amount SO2 (PBL) [DU] (05Jan2013 - 12Jan2013) US Dept of State Geographer Îmage Landsat © 2013 Google Jharkhan © 2013 Mapabc.com Madhya Pradesh lat 21.850338° lon 77.427513° elev 400 m Imagery Date: 4/9/2013 Eye **Output Files** sujarat ColumnAmountSO2_PBL.OMSO2e.003.AreaMap.2013-01-05.gif OMSO2e,003 Vertical Column Amount SO2 (PBL) [DU] (05Jan2013 - 12Jan2013) US Dept of State Geographer I.8 Chhattisgarh.... Odisha © 2013 Google © 2013 Mapabc.com Google earth Eye alt 1273.46 km M:latar 26:789419° lon 78.230331° elev 208 m

http://disc.sci.gsfc.nasa.gov/giovanni

January 5-12, 2013





SO2 NO2



Data/Images Access

Global Sulfur Dioxide Monitoring Homepage

- http://so2.gsfc.nasa.gov/
- Images, Documentation, Publications and Links

GES-DISC (Goddard Earth Science Data and Information Services Center)

- http://disc.sci.gsfc.nasa.gov/Aura/data-holdings/OMI/omso2e_v003.shtml
- Level 2, L2G, Level 3 (OMSO2e)
- HDF and NetCDF
- Documentation

GIOVANNI – An interactive visualization tool


- http://disc.sci.gsfc.nasa.gov/giovanni
- L2G, <u>Level 3 (OMSO2e)</u>
- HDF, ASCII, KML for Google Earth
- Subsetting available




Worldview

https://earthdata.nasa.gov/labs/worldview/

Uses Level 2 orbit data








NOTE: Data can be downloaded (HDF5 Format only) from Worldview BUT it does not subset the data so you don't know which orbit to pick for your region of interest, i.e. over CONUS.


Applied Remote SEnsing Training

Terra/MOPITT CO



CO plumes at an altitude of roughly 2 km (700 millibars) over the Houston area and extending out over the Bay of Galveston. This image represents a composite of data collected over a 10-day period, September 6-15, 2002, by MOPITT.

http://earthobservatory.nasa.gov/IOTD/view.php? id=3173






MOPITT- Measurements of Pollution In The Troposphere

HOMEPAGE = http://www.acd.ucar.edu/mopitt/

- Operational since March 2000
- A nadir sounding instrument
- Pixel resolution = 22km × 22km
- Swath width = 640 km
- Equator Crossing times = 10:00 AM, 10:10PM
- Vertical profiles of CO are obtained at 4.7 μm
 - Profile Units = Parts per Billion Volume mixing ratio (ppbv)
- Column CO are obtained at 2.2-2.4 μm
 - Column Units = molecules/cm2
- Current Version 6
- 9 vertical layers 900hPa 100hPa
- Level 3 gridded data are 1deg x1deg resolution
- Original data format is in HDF





Due to large gaps between swaths, it takes 3 days worth of data to create a global composite





MOPITT V6 Level 2 and 3 products include:

Homepage: http://www2.acd.ucar.edu/mopitt


A **TIR-only** product, similar to the MOPITT V4 product and V5 TIR-only product. *Example filename:* MOP02T-20010101-L2V16.2.1.he5.


A **NIR-only** product, similar to the MOPITT V5 NIR-only product. This dataset is produced only for daytime observations over land. <u>This product exhibits relatively large random errors and may require significant spatial and/or temporal averaging. *Example filename: MOP02N-20010101-L2V16.2.2.he5*.</u>


A **TIR/NIR** product, featuring the maximum sensitivity to near-surface CO. In this product, information from the NIR channels is exploited only in daytime observations over land. <u>This product exhibits relatively large random errors and may require significant spatial and/or temporal averaging. *Example filename: MOP02J-20010101-L2V16.2.3.he5*.</u>




MOPITT Product Images









NIR-only Daytime only
















Big MOPITT Limitation!! – COVERAGE

Due to large data gaps, do not expect to get daily coverage in your domain of interest!!



MOPITT Level 2, 3 Short names


Level 3 Level 2 Level 1 Level 0 Documentation			
Level Description: Derived Carbon Monoxide profiles			
Products	Temporal Resolution	Spatial Resolution	Temporal Coverage
MOP02 (HDF) MOPITT Derived CO Level 2 Data	Observations every 0.4 seconds.	Swath	03/03/2000 - Present
MOP02J (HDF) MOPITT Derived CO (Near and Thermal Infrared Radiances)	Observations every 0.4 seconds.	Swath	03/03/2000 - Present
MOP02N (HDF) MOPITT Derived CO (Near Infrared Radiances)	Observations every 0.4 seconds.	Swath	03/03/2000 - Present
MOP02T (HDF) MOPITT Derived CO (Thermal Infrared Radiances)	Observations every 0.4 seconds.	Swath	03/03/2000 - Present





File Names

AIRS














Homepage: http://www.acd.ucar.edu/mopitt/

MEASUREMENTS OF POLLUTION IN THE TROPOSPHERE (MOPITT)

THE MOPITT MISSION

MOPITT is an instrument flying on NASA's Earth Observing System Terra spacecraft, measuring tropospheric carbon monoxide (CO) on the global scale. MOPITT measurements enable scientists to analyze the distribution, transport, sources and sinks of CO, a trace gas produced by methane oxidation, fossil fuel consumption and biomass burning. MOPITT has been operational since March 2000.



MOPITT PRODUCT AVAILABILITY

MOPITT Version 6 Level 2 and Level 3 products are now available for the entire MOPITT mission. Generally, V6 Level 2 and Level 3 products should become available between three weeks and two months after the actual observations; this data latency is the result of the dependency on the MERRA reanalysis. Version 4 and Version 5 Level 2 and Level 3 products are also available. See Data Products for descriptions of the different products. User's Guides available on the Publications page should be consulted before attempting to analyze the MOPITT products.

Uses Level 3 1x1 degree gridded data with downloadable KMZ for Google Earth

MOPITT PRODUCT VISUALIZATION

Global plots of MOPITT CO products at 1 degree horizontal resolution can be viewed for each day and month of the MOPITT mission.

V6 Plots


- V6 TIR-only Daily Plots / Monthly Plots
- V6 NIR-only Daily Plots / Monthly Plots
- V6 TIR/NIR Daily Plots / Monthly Plots

V5 Plots

- V5 TIR-only Daily Plots / Monthly Plots
- . V5 NIR-only Daily Plots / Monthly Plots
- V5 TIR/NIR Daily Plots / Monthly Plots

V4 Plots

· V4 TIR-only Daily Plots / Monthly Plots





MOPITT Home » Visualization

VISUALIZATION

QUICK-LOOK IMAGES

Global plots of MOPITT CO products at 1 degree horizontal resolution can be viewed for each day and month of the MOPITT mission. See the News and Status page for dates of, and reasons for, missing data.

V6 Plots


- V6 TIR-only Daily Plots / Monthly Plots
- . V6 NIR-only Daily Plots / Monthly Plots
- . V6 TIR/NIR Daily Plots / Monthly Plots

V5 Plots

- . V5 TIR-only Daily Plots / Monthly Plots
- . V5 NIR-only Daily Plots / Monthly Plots
- . V5 TIR/NIR Daily Plots / Monthly Plots

V4 Plots

. V4 TIR-only Daily Plots / Monthly Plots



YEAR MONTH DAY LEVEL





PLOT

NEXT

MOPITT V5 TIR/NIR Daytime CO Retrievals - Quick-look Images










MOPITT now has an interactive visualizer

Homepage: http://www.acd.ucar.edu/mopitt/

MEASUREMENTS OF POLLUTION IN THE TROPOSPHERE (MOPITT)

THE MOPITT MISSION

MOPITT is an instrument flying on NASA's Earth Observing System Terra spacecraft, measuring tropospheric carbon monoxide (CO) on the global scale. MOPITT measurements enable scientists to analyze the distribution, transport, sources and sinks of CO, a trace gas produced by methane oxidation, fossil fuel consumption and biomass burning. MOPITT has been operational since March 2000.



MOPITT PRODUCT AVAILABILITY

MOPITT Version 6 Level 2 and Level 3 products are now available for the entire MOPITT mission. Generally, V6 Level 2 and Level 3 products should become available between three weeks and two months after the actual observations; this data latency is the result of the dependency on the MERRA reanalysis. Version 4 and Version 5 Level 2 and Level 3 products are also available. See Data Products for descriptions of the different products. User's Guides available on the Publications page should be consulted before attempting to analyze the MOPITT products.

- · NASA Data Archives via Reverb
- ASDC Data Pool
- ASDC MOPITT Subsetter (V5 Level 2 Products only)

MOPITT PRODUCT VISUALIZATION

Global plots of MOPITT CO products at 1 degree horizontal resolution can be viewed for each day and month of the MOPITT mission.

V6 Plots


- V6 TIR-only Daily Plots / Monthly Plots
- V6 NIR-only Daily Plots / Monthly Plots
- V6 TIR/NIR Daily Plots / Monthly Plots

V5 Plots

- V5 TIR-only Daily Plots / Monthly Plots
- . V5 NIR-only Daily Plots / Monthly Plots
- V5 TIR/NIR Daily Plots / Monthly Plots


V4 Plots

V4 TIR-only Daily Plots / Monthly Plots




Interactive Data Viewers for V4, V5, and V6 MOPITT products

MOPITT Interactive Data Viewer: http://mopfl.acd.ucar.edu:8080/webviewV6/




RetrievedCOSurfaceMixingRatio




Current Files:

- 1./MOPITT/V6T/Archive/L2/201301/0109/MOP02T-20130109-L2V16.2.1.he5
- 2. /MOPITT/V6T/Archive/L2/201301/0110/MOP02T-20130110-L2V16.2.1.he5
- 3./MOPITT/V6T/Archive/L2/201301/0111/MOP02T-20130111-L2V16.2.1.he5







MOPITT (+ all NASA EOS satellite) data are available at EOSDIS – NASA's Earth Observing System Data and Information System





Summary of Trace Gas Websites

GIOVANNI: http://disc.sci.gsfc.nasa.gov/giovanni

EOSDISC Reverb global search tool: http://reverb.echo.nasa.gov

GES-DISC: http://disc.sci.gsfc.nasa.gov

Mirador Archive: http://mirador.gsfc.nasa.gov

Worldview: https://earthdata.nasa.gov/labs/worldview/

Global Sulfur Dioxide Monitoring Homepage: http://so2.gsfc.nasa.gov/

Aura Homepage: http://aura.gsfc.nasa.gov

Aura Validation Data Center: http://avdc.gsfc.nasa.gov

OMSO2e Product: http://disc.sci.gsfc.nasa.gov/Aura/data-holdings/OMI/omso2e_v003.shtml

OMNO2d Product: http://disc.sci.gsfc.nasa.gov/Aura/data-holdings/OMI/omno2d_v003.shtml

MOPITT Homepage: http://www2.acd.ucar.edu/mopitt/

MOPITT Interactive Visualization tool:

http://mopfl.acd.ucar.edu:8080/webviewV6/selectmopittfile

